



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 45/06, 31/00, 31/10, 31/18, 38/13	A1	(11) International Publication Number: WO 97/29775 (43) International Publication Date: 21 August 1997 (21.08.97)
(21) International Application Number: PCT/US97/01422 (22) International Filing Date: 11 February 1997 (11.02.97) (30) Priority Data: 08/600,580 13 February 1996 (13.02.96) US (60) Parent Application or Grant (63) Related by Continuation US 08/600,580 (CON) Filed on 13 February 1996 (13.02.96) (71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GREGORY, Susan, A. [US/US]; 7269 Dartmouth, St. Louis, MO 63130 (US). ISAKSON, Peter, C. [US/US]; 2292 Ridgley Woods Drive, Clarkson Valley, MO 63005 (US). ANDERSON, Gary [US/US]; 1886 Woodhollow Drive #203, Maryland Heights, MO 63043 (US).	(74) Agents: BULOCK, Joseph, W. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: COMPOSITIONS COMPRISING A CYCLOOXYGENASE-2 INHIBITOR AND A LEUKOTRIENE B ₄ RECEPTOR ANTAGONIST (57) Abstract Treatment with a cyclooxygenase-2 inhibitor and a leukotriene B ₄ receptor antagonist is described as being useful in reducing recipient rejection of transplanted organs and for treatment of autoimmune diseases.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

COMPOSITIONS COMPRISING A CYCLOOXYGENASE-2 INHIBITOR AND A LEUKOTRIENE B₄ RECEPTOR ANTAGONIST

5

FIELD OF THE INVENTION

This invention is in the field of clinical immunology and relates to compositions having immunosuppressive properties. Of particular interest is a method of reducing recipient acute or chronic rejection of transplanted cells or organs, and for treatment of autoimmune diseases, hypersensitivity reactions of the acute or delayed type, allergic disorders, granulomas, meningitis, and septic shock by administering a cyclooxygenase-2 inhibitor and a leukotriene B₄ (LTB₄) receptor antagonist.

BACKGROUND OF THE INVENTION

Successful organ transplantation requires effective physiological and pharmacological intervention of the immune system of an organ recipient. Immunologic mechanisms are universal within the human species, but histocompatibility variations between organ donor and recipient may lead to rejection of donor tissue by stimulation of the recipient's immune system, except perhaps, in donor-recipient pairing of the monozygotic type. One approach to intervention of immune response in an organ transplant recipient, especially a recipient targeted for an allogenic graft, is by the use of immunosuppressive drugs. These drugs are used to prolong survival of transplanted organs in recipients in cases involving, for example, transplants of kidney, liver, heart, lung, bone marrow and pancreas.

There are several types of immunosuppressive drugs available for use in reducing organ rejection in transplantation. Such drugs fall within three major classes, namely: antiproliferative agents,

antiinflammatory-acting compounds and inhibitors of lymphocyte activation.

Examples of the class of cytotoxic or antiproliferative agents are azathioprine, cyclophosphamide and methotrexate. The compound azathioprine acts by interrupting DNA synthesis through inhibition of purine metabolism. The compound cyclophosphamide is an alkylating agent which interferes with enzyme actions and cell proliferation and interrupts DNA synthesis by binding to cellular DNA, RNA, and proteins. The compound methotrexate is a folic acid antagonist which interferes with nucleotide and protein synthesis. Drugs of the antiproliferative class may be effective immunosuppressives in patients with chronic inflammatory disorders and in organ transplant recipients by limiting cell proliferation. These drugs which abrogate mitosis and cell division have severe cytotoxic side effects on normal cell populations which have a high turn-over rate, such as bone marrow cells and cells of the gastrointestinal (GI) tract lining. Accordingly, such drugs often have severe side effects, particularly, lymphopenia, neutropenia, bone marrow depression, hemorrhagic cystitis, liver damage, increased incidence of malignancy, hair loss, GI tract disturbances, and infertility.

A second class of immunosuppressive drugs for use in transplantation is provided by compounds having antiinflammatory action. Representatives of this drug class are generally known as adrenal corticosteroids and have the advantage of not exerting globally systemic cytotoxic effects. These compounds usually act by preventing or inhibiting inflammatory responses or by reducing cytokine production, or by reducing chemotaxis, or by reducing neutrophil, macrophage or lymphocyte activation, or effector function. Typical examples of adrenal corticosteroids are prednisone and prednisolone which affect carbohydrate and protein metabolism as well as immune functions. Compounds of this class are

sometimes used in combination with cytotoxic agents, such as compounds of the antiproliferative class because the corticosteroids are significantly less toxic. But the adrenal corticosteroids lack specificity of effect and can exert a broad range of metabolic, antiinflammatory and immune effects. Typical side effects of this class include increased organ-recipient infections and interference with wound healing, as well as disturbing hemodynamic balance, carbohydrate and bone metabolism and mineral regulation.

A third class of immunosuppressive drugs for use in organ transplantation is provided by compounds which are immunomodulatory and generally prevent or inhibit leukocyte activation. Such compounds usually act by blocking activated T-cell effector functions or proliferation, or by inhibiting cytokine production, or by preventing or inhibiting activation, differentiation or effector functions of platelet, granulocyte, B-cell, or macrophage actions. The cyclosporin family of compounds is the leading example of drugs in this class. Such compounds are polypeptide fungal metabolites which have been found to be very effective in suppressing helper T-cells so as to reduce both cellular and humoral responses to newly-encountered antigens. Cyclosporins alter macrophage and lymphocyte activity by reducing cytokine production or secretion and, in particular, by interfering with activation of antigen-specific CD4 cells, by preventing IL-2 secretion and secretion of many T-cell products, as well as by interfering with expression of receptors for these lymphokines on various cell types. Cyclosporin A, in particular, has been used extensively as an immunosuppressive agent in organ transplantation. Other microbial metabolites include cyclosporins such as cyclosporin B and cyclosporin G, and another microbial product known as FK-506. Cyclosporin A suppresses humoral immunity as well as cell-mediated reactions. Cyclosporin A is indicated for organ rejection in kidney, liver, heart, pancreas, bone-

marrow and heart-lung transplants. Cyclosporin A is also useful in the treatment of autoimmune and inflammatory diseases, including rheumatoid arthritis, Crohn's disease, Graves' disease, severe psoriasis, aplastic anemia, multiple-sclerosis, alopecia areata, penphigus and penphigoid, dermatomyositis, polymyositis, Behcet's disease, uveitis, pulmonary sarcocidiosis, biliary cirrhosis, myasthenia gravis and atopic dermatitis.

10 Cyclosporins possess several significant disadvantages. While cyclosporins have provided significant benefits in organ transplantation, cyclosporins are non-specific immunosuppressives. Desirable immune reactions may be reduced against
15 foreign antigens. Tolerated dosages do not provide complete suppression of rejection response. Thus, immunologic reactions to transplanted tissue are not totally impeded, requiring concomitant treatment with prednisone, methylprednisolone, and/or other
20 immunosuppression agents, including monoclonal antibodies such as anti-CD3 or anti-CD5/CD7. Cyclosporins can produce severe side effects in many organ recipients, and show host-variable effects on the liver, kidney, the CNS and GI tract. Significant among
25 the adverse side effects are damage to the kidney and liver, hyperplasia of gum tissue, refractory hypertension and increased incidence of infections and malignancy.

Thus, the need remains for efficacious and
30 selective immunosuppressive drugs in organ transplantation, especially for grafts between less-than-perfectly matched donor-recipient pairs.

Prostaglandins and leukotrienes are lipid mediators produced in a variety of inflammatory disease states.
35 Both are products of metabolism of arachidonic acid. Cyclooxygenases (COX-1 and COX-2) are the enzymes that catalyze the conversion of arachidonic acid to prostaglandins. 5-Lipoxygenase (5-LO) catalyzes the

conversion of arachidonic acid to leukotrienes.

Products of both pathways have been described in association with transplant rejection in humans and animal models. Excess production of these mediators may
5 play a role in accelerating loss of the transplant function, particularly in the kidney. However, little research has been directed at determining direct effects of eicosanoids on tissue rejection.

Compounds which selectively inhibit cyclooxygenase-
10 2 have been described. U.S. patent 5,380,738 describes oxazoles which selectively inhibit cyclooxygenase-2. U.S. patent 5,344,991 describes cyclopentenones which selectively inhibit cyclooxygenase-2. U.S. patent 5,393,790 describes spiro compounds which selectively
15 inhibit cyclooxygenase-2. WO documents WO94/15932 describes thiophene and furan derivatives which selectively inhibit cyclooxygenase-2. WO94/27980 describes oxazoles which selectively inhibit cyclooxygenase-2. WO95/00501 describes compounds which
20 selectively inhibit cyclooxygenase-2. WO94/13635 describes compounds which selectively inhibit cyclooxygenase-2. WO94/20480 describes compounds which selectively inhibit cyclooxygenase-2. WO94/26731 describes compounds which selectively inhibit
25 cyclooxygenase-2. WO documents WO95/15316 describes pyrazolyl sulfonamide derivatives which selectively inhibit cyclooxygenase-2.

Compounds which affect leukotriene B₄ receptors have been described. U.S. Patent No. 5,384,318
30 describes substituted sulfonamides for the treatment of asthma. U.S. patent No. 5,246,965 describes aryl ethers as leukotriene B₄ receptor antagonists.

Combined therapies of NSAIDs and other reagents are known in the art. Combination analgesics have been
35 reported (W. Beaver, *Am. J. Med.*, 77, 38 (1984)) although such combinations do not substantially reduce adverse effects. The combination of NSAIDs and steroids have been described. A combination of indomethacin,

- steroid and lipopolysaccharide has been reported for the treatment of spinal injury (L. Guth et al., *Proc. Natl. Acad. Sci. USA*, **91**, 12308 (1994)). G. Hughes et al. describe combinations of corticosteroids with NSAIDs for the treatment of sunburn (*Dermatology*, **184**, 54 (1992)). C. Stewart et al. (*Clin. Pharmacol. Ther.*, **47**, 540 (1990)) describe the combination of naproxen and methotrexate as safe, although concurrent administrations of methotrexate with other NSAIDs have been reported to be toxic and sometimes fatal. A combination of a dual 5-lipoxygenase/cyclooxygenase inhibitor with a glucocorticoid is described for the treatment of skin disorders (K. Tramposch, *Inflammation*, **17**, 531 (1993)). Combinations of NSAIDs and steroids should be used in the treatment of scleritis only if patients are not responsive to any other treatment (S. Lightman and P. Watson, *Am. J. Ophthalmol.*, **108**, 95 (1989)). Combinations of cyclooxygenase inhibitors, lipoxygenase inhibitors, collagenase inhibitors and cytotoxic agents have been used in the treatment of non-small-cell lung cancers (B. Teicher et al., *Cancer. Chemother. Pharmacol.*, **33**, 515 (1994)). Combinations of naproxen with other NSAIDs have been described in the treatment of arthritis. R. Willikens and E. Segre (*Arthritis Rheum.*, **19**, 677 (1976)) describe the combination of aspirin and naproxen as being more effective than aspirin alone for the treatment of rheumatoid arthritis. Naproxen and acetaminophen together were described for treating the pain associated with arthritis (P. Seideman et al., *Acta Orthop. Scand.*, **64**, 285 (1993)). However, combinations of naproxen with indomethacin or ibuprofen offer no advantage in the treatment of arthritis (M. Seifert and C. Engler, *Curr. Med. Res. Opin.*, **7**, 38 (1980)).
- Tenidap has been described as inhibiting cyclooxygenases and cytokine-modifying [F. Breedveld, *Scand. J. Rheumatol.*, **23** (Supp. 100), 31 (1994)]. WO patent Publication 94/02448, published February 3, 1994,

describes hydroxamic acid derivatives as dual 5-lipoxygenase and cyclooxygenase inhibitors having immunosuppressant utility. U.S. Patent No. 4,595,699, to Terada et al., describes phenyl alkanoic acid derivatives as having analgesic, antiinflammatory and immune regulating activity. R. Bartlett et al. describe thiazolo(3,2-b)(1,2,4)triazin-7-ones as antiinflammatory agents with immunomodulating properties [*Drugs Exptl. Clin. Res.*, **15**, 521 (1989)]. J. Shaw and R. Greatorex [Adv. Prostaglandin, Thromboxane, Leukotriene Res., **13**, 219 (1985)] describe that whereas aspirin and sodium salicylate prolong graft survival, a cyclooxygenase inhibitor reduced the survival period. V. Fimiani, et al. describe some NSAID's that may have activity in the treatment of autoimmune diseases [*EOS-Revista di Immunologia and Immunofarmacologia*, **13**, 58 (1993)]. A. Badger et al. describe an indomethacin enhancement of suppressor cell population [*Immunopharm.*, **4**, 149 (1982)]. J. Shelby et al. [*Transplantation Proc.*, **19**, 1435 (1987)] describe indomethacin as reversing transfusion-induced graft prolongation. D. Latter et al. indicate that indomethacin was effective as an immunomodulator following burns [*J. Surg. Res.*, **43**, 246 (1987)]. J. Tarayre et al. describe indomethacin as having an effect in their delayed hypersensitivity models [*Arzneim.-Forsch./Drug Res.*, **40**, 1125 (1990)]. D. Braun et al indicate that a prostaglandin synthetase inhibitor may help prevent chemotherapy-induced decline in immune reactivity [*Proc. Am. Soc. Clin. Oncol.*, **4**, 21 Meeting, 223 (1985)]. Administration of tepoxalin (dual 5-LO and COX inhibitor) and cyclosporine has been described [Fung-Leung, et al., *Transplantation*, **60**, 362 (1995)] in suppression of graft versus host reaction although the effect of tepoxalin did not appear to be related to the inhibition of arachidonic acid metabolism.


There have been no reported combinations of a cyclooxygenase-2 selective inhibitor and a leukotriene

B₄ receptor antagonist as having a significant prolongation of graft survival.

DESCRIPTION OF THE INVENTION

Reduction in recipient rejection of a transplanted organ, or treatment of an autoimmune or inflammatory disease, or a hypersensitivity reaction of the acute or delayed type, an allergic reaction or asthmatic disorder, or treatment of dermatitis, arthritis, meningitis, granulomas, vasculitis, septic shock or graft vs. host response may be accomplished by a method to prevent or suppress immune responses in a recipient or treatment subject, which method comprises treating the subject with a therapeutically-effective amount of an immunosuppressive combination of a cyclooxygenase-2 inhibitor and a leukotriene B₄ receptor antagonist.

In addition, the invention describes a combination comprising a therapeutically-effective amount of a leukotriene B₄ receptor antagonist and a cyclooxygenase-2 inhibitor selected from Dupont Dup 697, Taisho NS-398, meloxicam, flosulide and compounds of Formula I



I

wherein A is a 5- or 6-member ring substituent selected from partially unsaturated or unsaturated heterocyclo and carbocyclic rings;

wherein R¹ is at least one substituent selected from heterocyclo, cycloalkyl, cycloalkenyl and aryl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxycarbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

wherein R² is selected from alkyl, and amino; and
wherein R³ is a radical selected from halo, alkyl,
alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl,
heterocyclooxy, alkyloxy, alkylthio, alkylcarbonyl,
5 cycloalkyl, aryl, haloalkyl, heterocyclo, cycloalkenyl,
aralkyl, heterocycloalkyl, acyl, alkylthioalkyl,
hydroxyalkyl, alkoxycarbonyl, arylcarbonyl,
aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl,
aryloxyalkyl, aralkylthioalkyl, aralkoxyalkyl,
10 alkoxyaralkoxyalkyl, alkoxycarbonylalkyl, aminocarbonyl,
aminocarbonylalkyl, alkylaminocarbonyl, N-
arylaminocarbonyl, N-alkyl-N-arylaminocarbonyl,
alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-
arylamino, N-aralkylamino, N-alkyl-N-aralkylamino, N-
15 alkyl-N-arylamino, aminoalkyl, alkylaminoalkyl, N-
arylaminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-
aralkylaminoalkyl, N-alkyl-N-arylaminominoalkyl, aryloxy,
aralkoxy, arylthio, aralkylthio, alkylsulfinyl,
alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-
20 arylaminosulfonyl, arylsulfonyl, N-alkyl-N-
arylaminosulfonyl; or a pharmaceutically-acceptable
salt thereof.

The invention would be useful for, but not limited
to, organ transplantation procedures and a variety of
25 disease states. For example, combinations of the
invention would be useful to treat a recipient of a
graft of a transplanted organ to reduce recipient
rejection of the graft or to reduce a donor leukocyte
response against the recipient's tissues. Such
30 combinations would be useful, in particular, for
transplants of bone marrow, kidney, liver, heart, heart-
lung and pancreas organs. Combinations of the invention
would also be useful in suppressing immune response in a
human or animal subject susceptible to or afflicted with
35 an autoimmune disease or inflammatory disease. Examples
of such treatable disease are graft vs. host disease,
systemic lupus erythematosus, multiple sclerosis,
myasthenia gravis, thyroiditis, Graves' disease,

autoimmune hemolytic anemia, aplastic anemia, autoimmune thrombocytopenia purpura, mixed connective tissue disease, idiopathic Addison's disease, Sjogren's syndrome, insulin dependent diabetes mellitus,

5 rheumatoid arthritis, osteoarthritis, skin and mucosal epithelial diseases such as psoriasis (in all its forms) lichen, chronic eczema, and pityriasis, glomerulonephritis, inflammatory bowel disease, Crohn's disease, alopecia areata, pemphigus and pemphigoid,

10 dermatomyositis, polymyositis, Behcet's disease, uveitis, pulmonary sarcoidosis, biliary cirrhosis, and atopic dermatitis. Combinations of the invention would also be useful in suppressing immune response in a human or animal subject susceptible to or afflicted with an

15 allergy, such as an asthmatic condition or reaction, urticaria or with airway hypersensitivity. The invention would also be useful in suppressing immune response in a human or animal subject afflicted with or susceptible to septic shock. Combinations of the

20 invention would also be useful in preventing or suppressing acute or delayed-type hypersensitivity responses or conditions resulting from or associated with hypersensitivity responses such as contact dermatitis, hemolytic anemias, antibody-induced

25 thrombocytopenia, Goodpasture's syndrome, hypersensitivity, pneumonitis, glomerulonephritis, granulomas, thyroiditis, encephelomyelitis, and meningitis. The invention would also be useful in the treatment of cancer, including leukemia, lymphoma and

30 solid tumors, including pancreatic, breast, colon, lung, epithelial and melanoma tumors.

Besides being useful for human treatment, these compounds are also useful for veterinary treatment of mammals, including companion animals and farm animals,

35 such as, but not limited to, horses, dogs, cats, cows, sheep and pigs.

Compositions of the invention would be useful in treating organs prior to transplant. For example, an

organ removed from a donor could be stored or transported in a bath containing an immunosuppressive composition of the invention. The immunosuppressive composition would act to inhibit donor leukocyte reactivity.

Compositions of the invention would also be useful in adjunct therapy involving, typically, coadministration with an additional immunosuppressive agent, such as a cyclosporin compound, or Fujisawa FK-506 (macrolide lactone) compound, or rapamycin, or a glucocorticoid, or an antiproliferative agent, or a monoclonal antibody such as an anti-CD3 (anti-T cell receptor antibody) or anti-CD5/CD7 or anti-CD4 agent, or an anti-IL-2 receptor (anti-cytokine receptor antibody) agent or an anti-IL-2 (anti-cytokine antibody), or Nippon NKT-01 (15-deoxyspergualin) or Syntex RS-61443.

The term "cyclooxygenase-2 inhibitor" embraces compounds which selectively inhibit cyclooxygenase-2 over cyclooxygenase-1. Preferably, the compounds have a cyclooxygenase-2 IC_{50} of less than about 0.5 μM , and also have a selectivity ratio of cyclooxygenase-2 inhibition over cyclooxygenase-1 inhibition of at least 50, and more preferably of at least 100. Even more preferably, the compounds have a cyclooxygenase-1 IC_{50} of greater than about 1 μM , and more preferably of greater than 20 μM . Such preferred selectivity may indicate an ability to reduce the incidence of common NSAID-induced side effects.

The term "leukotriene B_4 receptor antagonist" embraces compounds which selectively antagonize a leukotriene B_4 receptor with an IC_{50} of less than about 10 μM . More preferably, the leukotriene B_4 receptor antagonists have an IC_{50} of less than about 1 μM .

The phrase "combination therapy" (or "co-therapy"), in defining use of a cyclooxygenase-2 inhibitor agent and a leukotriene B_4 receptor antagonist agent, is intended to embrace administration of each agent in a

sequential manner in a regimen that will provide beneficial effects of the drug combination. The phrase also is intended to embrace co-administration of these agents in a substantially simultaneous manner, such as
5 in a single capsule having a fixed ratio of these active agents or in multiple, separate capsules for each agent.

The phrase "therapeutically-effective" is intended to qualify the amount of each agent for use in the combination therapy which will achieve the goal of
10 improvement in severity and the frequency of incidence over treatment of each agent by itself, while avoiding adverse side effects typically associated with alternative therapies.

Preferred leukotriene B₄ receptor antagonists
15 include calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, Terumo TMK-688, Boehringer Ingleheim BI-RM-270, Lilly LY 213024, Lilly LY 264086, Lilly LY 292728, Ono ONO LB457, Pfizer 105696, Perdue
20 Frederick PF 10042, Rhone-Poulenc Rorer RP 66153, SmithKline Beecham SB-201146, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Searle SC-53228, Sumitamo SM 15178, American Home Products WAY 121006, Bayer Bay-o-8276, Warner-Lambert CI-987, Warner-Lambert
25 CI-987BPC-15LY 223982, Lilly LY 233569, Lilly LY-255283, MacroNex MNX-160, Merck and Co. MK-591, Merck and CO. MK-886, Ono ONO-LB-448, Purdue Frederick PF-5901, Rhone-Poulenc Rorer RG 14893, Rhone-Poulenc Rorer RP 66364, Rhone-Poulenc Rorer RP 69698, Shionoogi S-2474, Searle
30 SC-41930, Searle SC-50505, Searle SC-51146, Searle SC-52798, SmithKline Beecham SK&F-104493, Leo Denmark SR-2566, Tanabe T-757 and Teijin TEI-1338.

More preferred leukotriene B₄ receptor antagonists include calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-
35 Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, Terumo TMK-688, Boehringer Ingleheim BI-RM-270, Lilly LY 213024, Lilly LY 264086, Lilly LY 292728, Ono ONO LB457, Pfizer 105696, Perdue

Frederick PF 10042, Rhone-Poulenc Rorer RP 66153, SmithKline Beecham SB-201146, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Searle SC-53228, Shionogi S-2472, Searle SC-52798, Leo Denmark SR-2566, 5 Tanabe T-757, Sumitamo SM 15178, and American Home Products WAY 121006.

Even more preferred leukotriene B₄ receptor antagonists include calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-10 615, Lilly LY-293111, Ono ONO-4057, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Pfizer 105696, and Terumo TMK-688.

A preferred class of compounds which inhibit cyclooxygenase-2 consists of compounds of Formula I 15 wherein A is selected from oxazolyl, isoxazolyl, thienyl, dihydrofuryl, furyl, pyrrolyl, pyrazolyl, thiazolyl, imidazolyl, isothiazolyl, cyclopentenyl, phenyl, and pyridyl; wherein R¹ is selected from 5- and 6-membered heterocyclo, lower cycloalkyl, lower 20 cycloalkenyl and aryl selected from phenyl, biphenyl and naphthyl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, cyano, carboxyl, lower alkoxy carbonyl, hydroxyl, lower 25 hydroxyalkyl, lower haloalkoxy, amino, lower alkylamino, phenylamino, nitro, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio; wherein R² is selected from lower alkyl and amino; and wherein R³ is a radical selected from halo, lower alkyl, 30 oxo, cyano, carboxyl, lower cyanoalkyl, heteroaryloxy, lower alkyloxy, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower hydroxylalkyl, lower aralkyl, acyl, phenylcarbonyl, lower alkoxyalkyl, heteroaryloxy, alkoxy carbonyl, 35 aminocarbonyl, alkylaminocarbonyl, alkylamino, aminoalkyl, alkylaminoalkyl, aryloxy, and aralkoxy; or a pharmaceutically-acceptable salt thereof.

A more preferred class of compounds which inhibit cyclooxygenase-2 consists of compounds of Formula I wherein A is selected from oxazolyl, isoxazolyl, dihydrofuryl, imidazolyl, and pyrazolyl; wherein R¹ is selected from 5- and 6-membered heterocyclo, lower cycloalkyl, lower cycloalkenyl and aryl selected from phenyl, biphenyl and naphthyl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, cyano, carboxyl, lower alkoxy carbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino, lower alkylamino, phenylamino, nitro, lower alkoxyalkyl, lower alkylsulfanyl, halo, lower alkoxy and lower alkylthio; wherein R² is amino; and wherein R³ is a radical selected from oxo, cyano, carboxyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, halo, lower alkyl, lower alkyloxy, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower hydroxyalkyl, lower aralkyl, acyl, phenylcarbonyl, lower alkoxyalkyl, 5- or 6-membered heteroaryloxy, aminocarbonyl, lower alkylaminocarbonyl, lower alkylamino, lower aminoalkyl, lower alkylaminoalkyl, phenyloxy, and lower aralkoxy; or a pharmaceutically-acceptable salt thereof.

An even more preferred class of compounds which inhibit cyclooxygenase-2 consists of compounds of Formula I wherein A is selected from oxazolyl, isoxazolyl, imidazolyl, and pyrazolyl; wherein R¹ is phenyl optionally substituted at a substitutable position with one or more radicals selected from methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, fluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, cyano, carboxyl, methoxycarbonyl, hydroxyl, hydroxymethyl, trifluoromethoxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, N,N-

- dipropylamino, N-butylamino, N-methyl-N-ethylamino, phenylamino, nitro, methoxymethyl, methylsulfinyl, fluoro, chloro, bromo, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and methylthio; wherein R² is amino; and wherein R³ is a radical selected from oxo, cyano, carboxyl, methoxycarbonyl, ethoxycarbonyl, carboxypropyl, carboxymethyl, carboxyethyl, cyanomethyl, fluoro, chloro, bromo, methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, fluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, cyclohexyl, phenyl, pyridyl, thienyl, thiazolyl, oxazolyl, furyl, pyrazinyl, hydroxymethyl, hydroxylpropyl, benzyl, formyl, phenylcarbonyl, methoxymethyl, furylmethyloxy, aminocarbonyl, N-methylaminocarbonyl, N,N-dimethylaminocarbonyl, N,N-dimethylamino, N-ethylamino, N,N-dipropylamino, N-butylamino, N-methyl-N-ethylamino, aminomethyl, N,N-dimethylaminomethyl, N-methyl-N-ethylaminomethyl, benzyloxy, and phenyloxy; or a pharmaceutically-acceptable salt thereof.

- A family of specific compounds of particular interest within Formula I consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 3-(3,4-difluorophenyl)-4-(4-methylsulfonylphenyl)-2-(5H)-furanone;
- 3-phenyl-4-(4-methylsulfonylphenyl)-2-(5H)-furanone;
- 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 3-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;

- 2-methyl-5-[1-[4-(methanysulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;
4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;
5 4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide;
4-[5-hydroxyethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;
[2-trifluoromethyl-5-(3,4-difluorophenyl)-4-oxazolyl]benzenesulfonamide;
10 4-[2-methyl-4-phenyl-5-oxazolyl]benzenesulfonamide; and
4-[5-(3-fluoro-4-methoxyphenyl-2-trifluoromethyl)-4-oxazolyl]benzenesulfonamide.

The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH₂-) radical. Where used, either alone or within other terms such as "haloalkyl", "alkylsulfonyl", "alkoxyalkyl" and "hydroxyalkyl", the term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about six carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl and the like. The term "alkenyl" embraces linear or branched radicals having at least one carbon-carbon double bond of two to about twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkenyl radicals are "lower alkenyl" radicals having two to about six carbon atoms. Examples of alkenyl radicals include ethenyl, propenyl, allyl, propenyl, butenyl and 4-methylbutenyl. The term "alkynyl" denotes linear or branched radicals having at least one carbon-carbon triple bond and having two to about twenty carbon atoms or, preferably, two to about

twelve carbon atoms. More preferred alkynyl radicals are "lower alkynyl" radicals having two to about ten carbon atoms. Most preferred are lower alkynyl radicals having two to about six carbon atoms. Examples of such radicals include propargyl, butynyl, and the like. The terms "alkenyl" and "lower alkenyl", embrace radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations. The term "cycloalkyl" embraces saturated carbocyclic radicals having three to about twelve carbon atoms. More preferred cycloalkyl radicals are "lower cycloalkyl" radicals having three to about eight carbon atoms. Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The term "cycloalkenyl" embraces partially unsaturated carbocyclic radicals having three to twelve carbon atoms. More preferred cycloalkenyl radicals are "lower cycloalkenyl" radicals having four to about eight carbon atoms. Examples of such radicals include cyclobutenyl, cyclopentenyl and cyclohexenyl. The term "halo" means halogens such as fluorine, chlorine, bromine or iodine. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. "Lower haloalkyl" embraces radicals having one to six carbon atoms. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of

which may be substituted with one or more hydroxyl radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms and one or more hydroxyl radicals. Examples of
5 such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl. The terms "alkoxy" and "alkyloxy" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals
10 are "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and tert-butoxy. The term "alkoxyalkyl" embraces alkyl radicals having one or more alkoxy radicals attached to the alkyl radical, that is,
15 to form monoalkoxyalkyl and dialkoxyalkyl radicals. The "alkoxy" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy radicals. More preferred haloalkoxy radicals are "lower haloalkoxy" radicals having one to
20 six carbon atoms and one or more halo radicals. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy. The term "aryl", alone or in combination, means a carbocyclic aromatic system
25 containing one, two or three rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane and biphenyl. Aryl moieties may also be substituted at a
30 substitutable position with one or more substituents selected independently from alkyl, alkoxyalkyl, alkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, alkoxy, aralkoxy, hydroxyl, amino, halo, nitro, alkylamino, acyl, cyano, carboxy,
35 aminocarbonyl, alkoxycarbonyl and aralkoxycarbonyl. The term "heterocyclo" embraces saturated, partially unsaturated and unsaturated heteroatom-containing ring-shaped radicals, where the heteroatoms may be selected

from nitrogen, sulfur and oxygen. Examples of saturated heterocyclo radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms (e.g. pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. morpholinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., thiazolidinyl, etc.). Examples of partially unsaturated heterocyclo radicals include dihydrothiophene, dihydropyran, dihydrofuran and dihydrothiazole. The term "heteroaryl" embraces unsaturated heterocyclo radicals. Examples of heteroaryl radicals include unsaturated 3 to 6 membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.) tetrazolyl (e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.), etc.; unsaturated condensed heterocyclo group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl (e.g., tetrazolo[1,5-b]pyridazinyl, etc.), etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.; unsaturated 3 to 6-membered heteromonocyclic group containing a sulfur atom, for example, thienyl, etc.; unsaturated 3- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.) etc.; unsaturated condensed heterocyclo group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. benzoxazolyl, benzoxadiazolyl, etc.); unsaturated 3 to 6-membered heteromonocyclic

group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl (e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.) etc.; unsaturated condensed

5 heterocyclo group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., benzothiazolyl, benzothiadiazolyl, etc.) and the like. The term "heteroaryl" also embraces radicals where heterocyclo radicals are fused with aryl radicals. Examples of such

10 fused bicyclic radicals include benzofuran, benzothiophene, and the like. Said "heterocyclo group" may have 1 to 3 substituents such as alkyl, hydroxyl, halo, alkoxy, oxo, amino and alkylamino. The term "alkylthio" embraces radicals containing a linear or

15 branched alkyl radical, of one to about ten carbon atoms attached to a divalent sulfur atom. More preferred alkylthio radicals are "lower alkylthio" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthio radicals are methylthio, ethylthio,

20 propylthio, butylthio and hexylthio. The term "alkylthioalkyl" embraces radicals containing an alkylthio radical attached through the divalent sulfur atom to an alkyl radical of one to about ten carbon atoms. More preferred alkylthioalkyl radicals are

25 "lower alkylthioalkyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthioalkyl radicals include methylthiomethyl. The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to about ten

30 carbon atoms, attached to a divalent $-S(=O)-$ radical. More preferred alkylsulfinyl radicals are "lower alkylsulfinyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylsulfinyl radicals include methylsulfinyl, ethylsulfinyl,

35 butylsulfinyl and hexylsulfinyl. The term "sulfonyl", whether used alone or linked to other terms such as "alkylsulfonyl", denotes a divalent radical, $-SO_2-$. "Alkylsulfonyl" embraces alkyl radicals attached to a

sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfonyl radicals include

5 methylsulfonyl, ethylsulfonyl and propylsulfonyl. The "alkylsulfonyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkylsulfonyl radicals. The terms "sulfamyl", "aminosulfonyl" and "sulfonamidyl" denote

10 $\text{NH}_2\text{O}_2\text{S}-$. The term "acyl" denotes a radical provided by the residue after removal of hydroxyl from an organic acid. Examples of such acyl radicals include alkanoyl and aroyl radicals. Examples of such lower alkanoyl radicals include formyl, acetyl, propionyl, butyryl,

15 isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, trifluoroacetyl. The term "carbonyl", whether used alone or with other terms, such as "alkoxycarbonyl", denotes $-(\text{C}=\text{O})-$. The term "aroyl" embraces aryl radicals with a carbonyl radical as defined above.

20 Examples of aroyl include benzoyl, naphthoyl, and the like and the aryl in said aroyl may be additionally substituted. The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes $-\text{CO}_2\text{H}$. The term "carboxyalkyl" embraces alkyl

25 radicals substituted with a carboxy radical. More preferred are "lower carboxyalkyl" which embrace lower alkyl radicals as defined above, and may be additionally substituted on the alkyl radical with halo. Examples of such lower carboxyalkyl radicals include carboxymethyl,

30 carboxyethyl and carboxypropyl. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl radical. More preferred are "lower alkoxycarbonyl" radicals with alkyl portions having one

35 to six carbons. Examples of such lower alkoxycarbonyl (ester) radicals include substituted or unsubstituted methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and hexyloxycarbonyl. The terms

"alkylcarbonyl", "arylcarbonyl" and "aralkylcarbonyl" include radicals having alkyl, aryl and aralkyl radicals, as defined herein, attached to a carbonyl radical. Examples of such radicals include substituted or unsubstituted methylcarbonyl, ethylcarbonyl, phenylcarbonyl and benzylcarbonyl. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, and diphenylethyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The terms benzyl and phenylmethyl are interchangeable. The term "heterocycloalkyl" embraces saturated and partially unsaturated heterocyclo-substituted alkyl radicals, such as pyrrolidinylmethyl, and heteroaryl-substituted alkyl radicals, such as pyridylmethyl, quinolylmethyl, thienylmethyl, furylethyl, and quinolylethyl. The heteroaryl in said heteroaralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The term "aralkoxy" embraces aralkyl radicals attached through an oxygen atom to other radicals. The term "aralkoxyalkyl" embraces aralkoxy radicals attached through an oxygen atom to an alkyl radical. The term "aralkylthio" embraces aralkyl radicals attached to a sulfur atom. The term "aralkylthioalkyl" embraces aralkylthio radicals attached through a sulfur atom to an alkyl radical. The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. More preferred are "lower aminoalkyl" radicals. Examples of such radicals include aminomethyl, aminoethyl, and the like. The term "alkylamino" denotes amino groups which are substituted with one or two alkyl radicals. Preferred are "lower alkylamino" radicals having alkyl portions having one to six carbon atoms. Suitable lower alkylamino may be monosubstituted N-alkylamino or disubstituted N,N-alkylamino, such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like. The term

"arylamino" denotes amino groups which are substituted with one or two aryl radicals, such as N-phenylamino. The "arylamino" radicals may be further substituted on the aryl ring portion of the radical. The term

5 "aralkylamino" embraces amino groups which are substituted with one or two aralkyl radicals. The terms "N-arylaminoalkyl" and "N-aryl-N-alkyl-aminoalkyl" denote aminoalkyl groups which are substituted with one aryl radical or one aryl and one alkyl radical,

10 respectively. Examples of such radicals include N-phenylaminomethyl and N-phenyl-N-methylaminomethyl. The term "aminocarbonyl" denotes an amide group of the formula $-C(=O)NH_2$. The term "alkylaminocarbonyl" denotes an aminocarbonyl group which has been

15 substituted with one or two alkyl radicals on the amino nitrogen atom. Preferred are "N-alkylaminocarbonyl" and "N,N-dialkylaminocarbonyl" radicals. More preferred are "lower N-alkylaminocarbonyl" and "lower N,N-dialkylaminocarbonyl" radicals with lower alkyl portions as defined above. The term "alkylaminoalkyl" embraces

20 radicals having one or more alkyl radicals attached to an aminoalkyl radical. The term "aryloxyalkyl" embraces radicals having an aryl radicals attached to an alkyl radical through a divalent oxygen atom. The term

25 "arylthioalkyl" embraces radicals having an aryl radicals attached to an alkyl radical through a divalent sulfur atom.

The present invention comprises a pharmaceutical composition comprising a therapeutically-effective

30 amount of a leukotriene B₄ receptor antagonist and a cyclooxygenase-2 inhibitor compound in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent.

The present invention also comprises a method of

35 treating immune-associated disorders in a subject, the method comprising treating the subject having or susceptible to such disorder with a therapeutically-effective amount of a leukotriene B₄ receptor antagonist

and a cyclooxygenase-2 inhibitor compound. The method of the present invention also includes prophylactic treatment.

Also included in the family of compounds of Formula I
5 are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is
10 pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and
15 phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclo, carboxylic and sulfonic classes of organic acids, example of which are


formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, p-hydroxybenzoic, 5 phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, cyclohexylaminosulfonic, algenic, β -hydroxybutyric, salicylic, galactaric and 10 galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, 15 chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate 20 acid or base with the compound of Formula I.

GENERAL SYNTHETIC PROCEDURES

25 The cyclooxygenase-2 inhibitor compounds of the invention can be synthesized according to the following procedures of Schemes I-X, wherein the R¹-R³ substituents are as defined for Formula I, above, except where further noted.

30

Scheme I




- Synthetic Scheme I shows the preparation of
- 5 cyclooxygenase-2 inhibitor compounds, as described in U.S. patent application Serial No. 08/223,629, which is incorporated by reference, embraced by Formula I.
- In step 1, ketone 1 is treated with a base, preferably NaOMe or NaH, and an ester, or ester equivalent, to
- 10 form the intermediate diketone 2 (in the enol form) which is used without further purification. In step 2, diketone 2 in an anhydrous protic solvent, such as absolute ethanol or acetic acid, is treated with the hydrochloride salt or the free base of a substituted
- 15 hydrazine at reflux to afford a mixture of pyrazoles 3 and 4. Recrystallization or chromatography affords 3 usually as a solid. Similar pyrazoles can be prepared by methods described in U.S. Pat. Nos. 4,146,721,

5,051,518, 5,134,142 and 4,914,121 which also are incorporated by reference.

Scheme II

5




Scheme II shows the four step procedure for forming cyclooxygenase-2 inhibitor pyrazoles 8 as described in U.S. patent application Serial No. 08/278,297 (where R^a is hydrido or alkyl) from ketones 5. In step 1, ketone 5 is reacted with a base, such as lithium bis(trimethylsilyl)amide or lithium diisopropylamide (LDA) to form the anion. In step 2, the anion is reacted with an acetylating reagent to provide diketone 6. In step 3, the reaction of diketone 6 with hydrazine or a substituted hydrazine, gives pyrazole 7. In step 4, the pyrazole 7 is oxidized with an oxidizing reagent, such as Oxone® (potassium peroxymonosulfate), 3-chloroperbenzoic acid (MCPBA) or hydrogen peroxide, to give a mixture of the desired 3-(alkylsulfonyl)phenyl-pyrazole 8 and the 5-(alkylsulfonyl)phenyl-pyrazole isomer. The desired pyrazole 8, usually a white or pale yellow solid, is obtained in pure form either by chromatography or recrystallization.

Alternatively, diketone 6 can be formed from ketone 5 by treatment with a base, such as sodium

hydride, in a solvent, such as dimethylformamide, and further reacting with a nitrile to form an aminoketone. Treatment of the aminoketone with acid forms the diketone 6. Similar pyrazoles can be
5 prepared by methods described in U.S. Pat. No. 3,984,431 which is incorporated by reference.

Scheme III




- 5 Cyclooxygenase-2 inhibitor diaryl/heteroaryl thiophenes (where T is S, and R^b is alkyl) can be prepared by the methods described in U.S. Patent Nos. 4,427,693, 4,302,461, 4,381,311, 4,590,205, and 4,820,827, and PCT documents WO 95/00501 and
- 10 WO94/15932, which are incorporated by reference. Similar pyrroles (where T is N), furanones and furans

(where T is O) can be prepared by methods described in PCT documents WO 95/00501 and WO94/15932.


Scheme IV

5




Cyclooxygenase-2 inhibitor diaryl/heteroaryl oxazoles can be prepared by the methods described in U.S. Patent Nos. 3,743,656, 3,644,499 and 3,647,858, and PCT documents WO 95/00501 and WO94/27980, which are incorporated by reference.

Scheme V



- 5 Cyclooxygenase-2 inhibitor diaryl/heteroaryl isoxazoles can be prepared by the methods described in United States application Serial No. 08/387,680, PCT documents WO92/05162, and WO92/19604, and European Publication EP 26928 which are incorporated by
- 10 reference. Sulfonamides 24 can be formed from the hydrated isoxazole 23 in a two step procedure. First, hydrated isoxazole 23 is treated at about 0 °C with two or three equivalents of chlorosulfonic acid to form the corresponding sulfonyl chloride. In step
- 15 two, the sulfonyl chloride thus formed is treated with concentrated ammonia to provide the sulfonamide derivative 24.

Scheme VI




Scheme VI shows the three step preparation of the
 5 cyclooxygenase-2 inhibitor imidazoles 29 of the
 present invention. In step 1, the reaction of
 substituted nitriles (R^1CN) 25 with primary
 phenylamines 26 in the presence of alkylaluminum
 reagents such as trimethylaluminum, triethylaluminum,
 10 dimethylaluminum chloride, diethylaluminum chloride in
 the presence of inert solvents such as toluene,
 benzene, and xylene, gives amidines 27. In step 2,
 the reaction of amidine 27 with 2-haloketones (where X
 is Br or Cl) in the presence of bases, such as sodium
 15 bicarbonate, potassium carbonate, sodium carbonate,
 potassium bicarbonate or hindered tertiary amines such
 as *N,N'*-diisopropylethylamine, gives the 4,5-

dihydroimidazoles 28 (where R^b is alkyl). Some of the suitable solvents for this reaction are isopropanol, acetone and dimethylformamide. The reaction may be carried out at temperatures of about 20°C to about 90°C. In step 3, the 4,5-dihydroimidazoles 28 may be dehydrated in the presence of an acid catalyst such as 4-toluenesulfonic acid or mineral acids to form the 1,2-disubstituted imidazoles 29 of the invention. Suitable solvents for this dehydration step are e.g., toluene, xylene and benzene. Trifluoroacetic acid can be used as solvent and catalyst for this dehydration step.

In some cases (e.g., where R^3 = methyl or phenyl) the intermediate 28 may not be readily isolable. The reaction, under the conditions described above, proceeds to give the targeted imidazoles directly.

Similarly, imidazoles can be prepared having the sulfonylphenyl moiety attached at position 2 and R^1 attached at the nitrogen atom at position 1. Diaryl/heteroaryl imidazoles can be prepared by the methods described in U.S. Patent Nos. 4,822,805, U.S. application Serial No. 08/282,395 and PCT document WO 93/14082, which are incorporated by reference.

Scheme VII



- 5 The subject imidazole cyclooxygenase-2 inhibitor compounds 36 of this invention may be synthesized according to the sequence outlined in Scheme VII. Aldehyde 30 may be converted to the protected cyanohydrin 31 by reaction with a trialkylsilyl
- 10 cyanide, such as trimethylsilyl cyanide (TMSCN) in the presence of a catalyst such as zinc iodide (ZnI₂) or

potassium cyanide (KCN). Reaction of cyanohydrin 31 with a strong base followed by treatment with benzaldehyde 32 (where R^2 is alkyl) and using both acid and base treatments, in that order, on workup gives benzoin 33. Examples of strong bases suitable for this reaction are lithium diisopropylamide (LDA) and lithium hexamethyldisilazane. Benzoin 33 may be converted to benzil 34 by reaction with a suitable oxidizing agent, such as bismuth oxide or manganese dioxide, or by a Swern oxidation using dimethyl sulfoxide (DMSO) and trifluoroacetic anhydride. Benzil 34 may be obtained directly by reaction of the anion of cyanohydrin 31 with a substituted benzoic acid halide. Any of compounds 33 and 34 may be used as intermediates for conversion to imidazoles 35 (where R^2 is alkyl) according to chemical procedures known by those skilled in the art and described by M. R. Grimmett, "Advances in Imidazole Chemistry" in *Advances in Heterocyclic Chemistry*, 12, 104 (1970). The conversion of 34 to imidazoles 35 is carried out by reaction with ammonium acetate and an appropriate aldehyde (R^3CHO) in acetic acid. Benzoin 36 may be converted to imidazoles 38 by reaction with formamide. In addition, benzoin 36 may be converted to imidazoles by first acylating with an appropriate acyl group (R^3CO-) and then treating with ammonium hydroxide. Those skilled in the art will recognize that the oxidation of the sulfide (where R^2 is methyl) to the sulfone may be carried out at any point along the way beginning with compounds 35, and including oxidation of imidazoles 38, using, for examples, reagents such as hydrogen peroxide in acetic acid, *m*-chloroperoxybenzoic acid (MCPBA) and potassium peroxymonosulfate (OXONE®).


35

Diaryl/heteroaryl imidazoles can be prepared by the methods described in U.S. Patent Nos. 3,707,475, 4,686,231, 4,503,065, 4,472,422,

4,372,964, 4,576,958, 3,901,908, U.S. application
Serial No. 08/281,903 European publication EP 372,445,
and PCT document WO 95/00501, which are incorporated
by reference.

5


Scheme VIII



10


Diaryl/heteroaryl cyclopentene cyclooxygenase-2
inhibitors can be prepared by the methods described in
U.S. Patent No. 5,344,991, and PCT document WO
95/00501, which are incorporated by reference.

Scheme IX



- 5 Similarly, Synthetic Scheme IX shows the
 procedure for the preparation of 1,2-diarylbenzene
 cyclooxygenase-2 inhibitor agents 44 from 2-bromo-
 biphenyl intermediates 43 (prepared similar to that
 described in Synthetic Scheme VIII) and the
 10 appropriate substituted phenylboronic acids. Using a
 coupling procedure similar to the one developed by
 Suzuki et al. [*Synth. Commun.*, 11, 513 (1981)],
 intermediates 43 are reacted with the boronic acids in
 toluene/ethanol at reflux in the presence of a Pd⁰
 15 catalyst, e.g.,
 tetrakis(triphenylphosphine)palladium(0), and 2M
 sodium carbonate to give the corresponding 1,2-
 diarylbenzene antiinflammatory agents 44 of this
 invention. Such terphenyl compounds can be prepared
 20 by the methods described in U.S. application Serial
 No. 08/346,433, which is incorporated by reference.

Scheme X



Diaryl/heteroaryl thiazole cyclooxygenase-2

5 inhibitors can be prepared by the methods described in U.S. Patent No. 4,051,250, 4,632,930, U.S. application Serial No. 08/281,288, European Application EP 592,664, and PCT document WO 95/00501, which are incorporated by reference. Isothiazoles can be prepared as described in PCT document WO 95/00501.

Diaryl/heteroaryl pyridine cyclooxygenase-2

15 inhibitors can be prepared by the methods described in U.S. Patent Nos. 5,169,857, 4,011,328, 4,533,666, U.S. application Serial No. 08/386,843 and U.S. application Serial No. 08/387,150 which are incorporated by reference.

The following examples contain detailed descriptions of the methods of preparation of combinations with compounds of Formula I. These detailed descriptions fall within the scope, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These detailed descriptions are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated. All compounds showed NMR spectra consistent with their assigned structures.

Example 1

4-[5-(4-Chlorophenyl)-3-(trifluoromethyl)-
1H-pyrazol-1-yl]benzenesulfonamide

5 Step 1: Preparation of 4,4,4-trifluoro-1-[4-(chloro)phenyl]-butane-1,3-dione.

Ethyl trifluoroacetate (23.52 g, 166 mmol) was dissolved in methyl tert-butyl ether (75 mL). To the stirred solution was added 25 weight % sodium methoxide
10 (40 mL, 177 mmol). 4'-Chloroacetophenone (23.21 g, 150 mmol) was dissolved in methyl tert-butyl ether (20 mL) and added to the reaction dropwise. After stirring overnight (15.75 hours), 3N HCl (70 mL) was added. The organic layer was collected, washed with brine (75 mL),
15 dried over MgSO₄, filtered, and concentrated in vacuo to give a yellow-orange solid. The solid was recrystallized from isooctane to give the dione (31.96 g, 85%): mp 66-67°C.

20 Step 2: Preparation of 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide.

4-Sulphonamidophenyl hydrazine hydrochloride (982 mg, 4.4 mmol, 1.1 equiv.) was added to a stirred solution of 4,4,4-trifluoro-1-[4-(chloro)phenyl]-butane-
25 1,3-dione from Step 1 (1.00 g, 4.0 mmol) in ethanol (50 mL). The reaction was heated to reflux and stirred for 20 hours. After cooling to room temperature, the reaction mixture was concentrated in vacuo. The residue was taken up in ethyl acetate and washed with water and
30 brine. The residue was dried over MgSO₄, filtered, and concentrated in vacuo to give a light brown solid. The solid was recrystallized from ethyl acetate and isooctane to give the pyrazole (1.28 g, 80%): mp 143-145°C; EI GC-MS M⁺ = 401.

35

Example 2

4-[5-(3-Fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide

Step 1: Preparation of 3'-fluoro-4'-methoxy-acetophenone.

Acetyl chloride (51.0 g, 0.65 mol) was added dropwise to a stirred solution of aluminum chloride (80.0 g, 0.6 mol) and chloroform (750 mL), maintaining the temperature between 5-10°C. The mixture was stirred for 10 minutes at 5°C before the dropwise addition of 2-fluoroanisole (62.6 g, 0.5 mol). The mixture was stirred at 0-10°C for 1 hour and poured into ice (1 L). The resultant layers were separated and the aqueous layer was extracted with dichloromethane (2x250 mL). The combined organic layers were washed with water (2x150 mL), dried over anhydrous MgSO₄, filtered and concentrated in vacuo to a volume of 300 mL. Hexanes were added and a white solid formed which was isolated by filtration and air dried. This material was recrystallized from a mixture of dichloromethane and hexanes to afford material suitable for use in the next step (77.2 g, 92%): mp 92-94°C.

Step 2: Preparation of 4,4-difluoro-1-(3-fluoro-4-methoxyphenyl)-butane-1,3-dione.

Ethyl difluoroacetate (4.06 g, 32.7 mmol) was dissolved in methyl tert-butyl ether (50 mL). To the stirred solution was added 25 weight % sodium methoxide (7.07 g, 32.7 mmol) followed by 3'-fluoro-4'-methoxyacetophenone from Step 1 (5.0 g, 29.7 mmol). After stirring for 16 hours, 1N HCl (50 mL) was added. The organic layer was collected and washed with water (2x50 mL), dried over anhydrous MgSO₄, filtered, and added to hexanes to precipitate a tan solid (7.0 g, 96%): mp 70-72°C.


Step 3: Preparation of 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide.

4,4-Difluoro-1-(3-fluoro-4-methoxyphenyl)-butane-1,3-dione from Step 2 (7.0 g, 28.4 mmol) was dissolved in ethanol (150 mL). To the stirred mixture was added

4-sulphonamidophenyl hydrazine hydrochloride (7.4 g, 33 mmol) and stirred at reflux overnight (16 hours). The mixture was cooled and water was added until crystals slowly appeared. The product was isolated by filtration and air dried to provide the desired product as a light tan solid (9.8 g, 87%): mp 159-161°C. Anal. Calc'd. for $C_{17}H_{14}N_3SO_3F_3$: C, 51.38; H, 3.55; N, 10.57. Found: C, 51.46; H, 3.52; N, 10.63.

10

Example 3



7-[3-[2-(Cyclopropylmethyl)-3-methoxy-4-
15 [(methylamino)carbonyl]phenoxy]propoxy]-3,4-dihydro-8-propyl-
2H-1-benzopyran-2-propanoic acid is prepared as in U.S.
Patent No. 5,310,951, which is incorporated by reference.

20

BIOLOGICAL EVALUATION

A combination of a cyclooxygenase-2 inhibitor and a leukotriene B₄ receptor antagonist is evaluated as described in the following tests.

25

Transplantation and Evaluation of Graft Rejection

The method of skin grafting used has been previously described [D. Steinmuller, Skin Grafting. Surgical Techniques in Immunology, Methods Enzymol.
30 108, 20 (1984)]. Briefly, a tailskin from an 8-12 week old male B10.Br mouse is removed and stored in cold saline. Male C57BL/10 mice are anesthetized, and

their backs are shaved. The backs are scrubbed with alcohol, and a 1 cm² piece of skin is removed. A similar size piece of skin is cut from the tailskin of the B10.Br mouse and placed in the excised area on the C57BL/10 animal's back. A petroleum jelly coated bandage is placed over the graft and held in place by a bandage. Compounds are prepared as a suspension in 0.5% methylcellulose (Sigma, St. Louis, MO), and 0.025% Tween[®] 20 (Sigma). The compounds are administered by i.p. injection in a volume of 0.1 ml beginning on the day of skin grafting and continuing until transplant rejection. Cyclosporin A (csa) is purchased as "Sandimmune Injection" at a pharmacy. Compounds are administered alone or as combinations of a COX-2 and a leukotriene B₄ receptor antagonist. Bandages are left in place until 8 days post grafting. At that time they are removed, and the grafts are observed daily for signs of rejection. Rejection is determined by complete blackening or scabbing of the grafted skin. The animals are dosed at one of the following dosing ranges:

Example 1 @ M,W,F @ 10 mpk/day;
Example 2 @ 30 mpk/day, q.d.;
Example 3 @ 10 mpk/day, q.d.;
csa @ 5 mpk/day, b.d.

The combinations of a COX-2 inhibitor or the leukotriene B₄ receptor antagonist should be active in delaying graft rejection at a dosage of about 10-20 mg per kg body weight. The coadministration of a COX-2 inhibitor or the leukotriene B₄ receptor antagonist with a low dose of the immunosuppressant Cyclosporin A should enhance prolongation of graft survival and may have additive or synergistic effects when combined with cyclosporin.

Example 4

A formulation is prepared having the following components:

700 mg of a cyclooxygenase-2 inhibitor and 700 mg of a leukotriene B₄ receptor antagonist.

Example 5

A formulation is prepared having the following components:

350 mg of 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide and 700 mg of 7-[3-[2-(cyclopropylmethyl)-3-methoxy-4-[(methylamino)carbonyl]phenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoic acid.

Also embraced within this invention is a class of pharmaceutical compositions comprising the active compounds of this combination therapy in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The active compounds and composition may, for example, be administered orally, intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier.

The amount of therapeutically active compounds that are administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the disease, the route and frequency of administration, and the particular compound employed, and thus may vary widely. The pharmaceutical compositions may contain active ingredients in the range of about 0.1 to 2000 mg, preferably in the range of about 0.5 to 500 mg and most preferably between about 1 and 100 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between about 0.05 and about 20 mg/kg body weight and most preferably between about 0.1 to 10 mg/kg body weight, may be appropriate. The daily dose can be administered in one to four doses per day.

In the case of psoriasis and other skin conditions, it may be preferable to apply a topical preparation of compounds of this invention to the affected area two to four times a day.

For disorders of the eye or other external tissues, e.g., mouth and skin, the formulations are preferably applied as a topical ointment or cream, or as a suppository, containing the active ingredients in a total amount of, for example, 0.075 to 30% w/w, preferably 0.2 to 20% w/w and most preferably 0.4 to 15% w/w. When formulated in an ointment, the active ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of

such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be
5 accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in
10 contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the encapsulating agent may also function
15 as the membrane.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one
20 emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with
25 or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers
30 suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, and sodium lauryl sulfate, among others.

The choice of suitable oils or fats for the
35 formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream

should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredients are dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The antiinflammatory active ingredients are preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% and particularly about 1.5% w/w.

For therapeutic purposes, the active compounds of this combination invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or

suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds
5 may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the
10 pharmaceutical art.


Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

15

What is claimed is :

1. A method to suppress immune, acute or delayed-type hypersensitivity response in a subject, said
 5 method comprising treating the subject with a therapeutically-effective amount of a leukotriene B₄ receptor antagonist and a cyclooxygenase-2 inhibitor selected from Dupont Dup 697, Taisho NS-398, meloxicam, flosulide and compounds of Formula I

10



I

- wherein A is a 5- or 6-member ring substituent selected from partially unsaturated or unsaturated
 15 heterocyclo and carbocyclic rings;

- wherein R¹ is at least one substituent selected from heterocyclo, cycloalkyl, cycloalkenyl and aryl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxy, carbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

- wherein R² is selected from alkyl, and amino; and

- 25 wherein R³ is a radical selected from halo, alkyl, alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl, heterocycloalkoxy, alkoxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, heterocyclo, cycloalkenyl, aralkyl, heterocycloalkyl, acyl, alkylthioalkyl, hydroxyalkyl, alkoxy, carbonyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl, arylalkoxyalkyl, aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl, alkoxy, carbonylalkyl, aminocarbonyl, aminocarbonylalkyl, alkylaminocarbonyl, N-arylaminocarbonyl, N-alkyl-N-arylaminocarbonyl, alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-

arylamino, N-aralkylamino, N-alkyl-N-aralkylamino, N-alkyl-N-arylamino, aminoalkyl, alkylaminoalkyl, N-arylaminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-aralkylaminoalkyl, N-alkyl-N-arylaminoalkyl, aryloxy,
5 aralkoxy, arylthio, aralkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-arylaminosulfonyl, arylsulfonyl, N-alkyl-N-arylaminosulfonyl;

or a pharmaceutically-acceptable salt thereof.

10

2. The method of Claim 1 wherein said leukotriene B₄ receptor antagonist and said cyclooxygenase-2 inhibitor are administered in a sequential manner.

15

3. The method of Claim 1 wherein said leukotriene B₄ receptor antagonist and said cyclooxygenase-2 inhibitor are administered in a substantially simultaneous manner.

20

4. The method of Claim 1 wherein the leukotriene B₄ receptor antagonist is selected from calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, Terumo TMK-688, Boehringer Ingelheim BI-RM-270, Lilly LY 213024, Lilly LY 264086, Lilly LY 292728, Ono ONO LB457, Pfizer 105696, Perdue Frederick PF 10042, Rhone-Poulenc Rorer RP 66153, SmithKline Beecham SB-201146, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Searle SC-53228,
25 Sumitamo SM 15178, American Home Products WAY 121006, Bayer Bay-o-8276, Warner-Lambert CI-987, Warner-Lambert CI-987BPC-15LY 223982, Lilly LY 233569, Lilly LY-255283, MacroNex MNX-160, Merck and Co. MK-591, Merck and CO. MK-886, Ono ONO-LB-448, Purdue Frederick
30 PF-5901, Rhone-Poulenc Rorer RG 14893, Rhone-Poulenc Rorer RP 66364, Rhone-Poulenc Rorer RP 69698, Shionogi S-2474, Searle SC-41930, Searle SC-50505, Searle SC-51146, Searle SC-52798, SmithKline Beecham

SK&F-104493, Leo Denmark SR-2566, Tanabe T-757 and Teijin TEI-1338.

5. The method of Claim 4 wherein the leukotriene
5 B₄ receptor antagonist is selected from calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, Terumo TMK-688, Boehringer Ingleheim BI-RM-270, Lilly LY 213024, Lilly LY 264086, Lilly LY 292728, Ono
10 ONO LB457, Pfizer 105696, Perdue Frederick PF 10042, Rhone-Poulenc Rorer RP 66153, SmithKline Beecham SB-201146, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Searle SC-53228, Shionogi S-2472, Searle SC-52798, Leo Denmark SR-2566, Tanabe T-757,
15 Sumitamo SM 15178, and American Home Products WAY 121006.

6. The combination of Claim 5 wherein the
leukotriene B₄ receptor antagonist is selected from
20 calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Pfizer 105696, and Terumo TMK-688.

25

7. The method of Claim 1 wherein A is selected from oxazolyl, isoxazolyl, dihydrofuryl, imidazolyl, and pyrazolyl; wherein R¹ is selected from 5- and 6-membered heterocyclo, and aryl selected from phenyl,
30 biphenyl and naphthyl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, cyano, carboxyl, lower alkoxy, carbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino,
35 lower alkylamino, phenylamino, nitro, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio; wherein R² is amino; and wherein R³ is a radical selected from oxo, cyano, carboxyl, lower

alkoxycarbonyl, lower carboxyalkyl, lower cyanoalkyl, halo, lower alkyl, lower alkyloxy, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower hydroxylalkyl, lower aralkyl, acyl,
5 phenylcarbonyl, lower alkoxyalkyl, 5- or 6-membered heteroaryloxy, aminocarbonyl, lower alkylaminocarbonyl, lower alkylamino, lower aminoalkyl, lower alkylaminoalkyl, phenyloxy, and lower aralkoxy; or a pharmaceutically-acceptable salt thereof.

10

8. The method of Claim 7 wherein A is selected from oxazolyl, isoxazolyl, imidazolyl, and pyrazolyl; wherein R¹ is phenyl optionally substituted at a substitutable position with one or more radicals
15 selected from methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, trifluoromethyl, cyano, carboxyl, methoxycarbonyl, hydroxyl, hydroxymethyl, trifluoromethoxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, N,N-dipropylamino, N-butylamino, N-methyl-N-ethylamino, nitro,
20 methoxymethyl, methylsulfinyl, fluoro, chloro, bromo, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and methylthio; wherein R² is amino; and wherein R³ is a radical selected from oxo, cyano, carboxyl, methoxycarbonyl, ethoxycarbonyl, carboxypropyl, carboxymethyl, carboxyethyl, cyanomethyl, fluoro, chloro, bromo, methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl,
25 dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, fluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, cyclohexyl, phenyl, pyridyl, thienyl, thiazolyl, oxazolyl, furyl,
30 pyrazinyl, hydroxylmethyl, hydroxylpropyl, benzyl, formyl, phenylcarbonyl, methoxymethyl, furylmethyloxy, aminocarbonyl, N-methylaminocarbonyl, N,N-dimethylaminocarbonyl, N,N-dimethylamino, N-ethylamino,

N,N-dipropylamino, N-butylamino, N-methyl-N-ethylamino, aminomethyl, N,N-dimethylaminomethyl, N-methyl-N-ethylaminomethyl, benzyloxy, and phenyloxy; or a pharmaceutically-acceptable salt thereof.

5

9. The method of Claim 8 wherein the cyclooxygenase-2 inhibitor is selected from compounds, their prodrugs and their pharmaceutically-acceptable salts, of the group consisting of

10

3-(3,4-difluorophenyl)-4-(4-methylsulfonylphenyl)-2-(5H)-furanone;

3-phenyl-4-(4-methylsulfonylphenyl)-2-(5H)-furanone;

15 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

20 3-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;

2-methyl-5-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;

25 4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;

4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide;

4-[5-hydroxyethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;

30 [2-trifluoromethyl-5-(3,4-difluorophenyl)-4-oxazolyl]benzenesulfonamide;

4-[2-methyl-4-phenyl-5-oxazolyl]benzenesulfonamide; and


4-[5-(3-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-oxazolyl]benzenesulfonamide.

35

10. A combination comprising a therapeutically-effective amount of a cyclooxygenase-2 inhibitor, a leukotriene B₄ receptor antagonist and an immunosuppressive drug selected from antiproliferative

agents, antiinflammatory-acting compounds and inhibitors of leukocyte activation.

11. The combination of Claim 10 wherein the
 5 cyclooxygenase-2 inhibitor is selected from Dupont Dup-697, Taisho NS-398, meloxicam, flosulide and compounds of Formula I



10

wherein A is a 5- or 6-member ring substituent selected from partially unsaturated or unsaturated heterocyclo and carbocyclic rings;

- wherein R¹ is at least one substituent selected
 15 from heterocyclo, cycloalkyl, cycloalkenyl and aryl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxycarbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino,
 20 nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

wherein R² is selected from alkyl, and amino; and

- wherein R³ is a radical selected from halo, alkyl, alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl,
 25 heterocycloxy, alkyloxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, heterocyclo, cycloalkenyl, aralkyl, heterocycloalkyl, acyl, alkylthioalkyl, hydroxyalkyl, alkoxycarbonyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl,
 30 aryloxyalkyl, aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl, alkoxycarbonylalkyl, aminocarbonyl, aminocarbonylalkyl, alkylaminocarbonyl, N-arylaminocarbonyl, N-alkyl-N-arylaminocarbonyl, alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-
 35 arylamino, N-aralkylamino, N-alkyl-N-aralkylamino, N-

alkyl-N-arylamino, aminoalkyl, alkylaminoalkyl, N-arylaminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-aralkylaminoalkyl, N-alkyl-N-arylaminoalkyl, aryloxy, aralkoxy, arylthio, aralkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-arylaminosulfonyl, arylsulfonyl, N-alkyl-N-arylaminosulfonyl;

or a pharmaceutically-acceptable salt thereof.

10 12. The combination of Claim 10 wherein the leukotriene B₄ receptor antagonist is selected from, calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, Terumo TMK-688, Boehringer
15 Ingleheim BI-RM-270, Lilly LY 213024, Lilly LY 264086, Lilly LY 292728, Ono ONO LB457, Pfizer 105696, Perdue Frederick PF 10042, Rhone-Poulenc Rorer RP 66153, SmithKline Beecham SB-201146, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Searle SC-53228,
20 Shionogi S-2472, Searle SC-52798, Leo Denmark SR-2566, Tanabe T-757, Sumitamo SM 15178, and American Home Products WAY 121006.

25 13. The combination of Claim 12 wherein the leukotriene B₄ receptor antagonist is selected from calcitriol, ontazolast, Bayer Bay-x-1005, Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Warner-Lambert BPC-15,
30 Pfizer 105696, Shionogi S-2472, Searle SC-52798, Leo Denmark SR-2566, Tanabe T-757 and Terumo TMK-688.

35 14. The combination of Claim 11 wherein A is selected from oxazolyl, isoxazolyl, thienyl, dihydrofuryl, furyl, pyrrolyl, pyrazolyl, thiazolyl, imidazolyl, isothiazolyl, cyclopentenyl, phenyl, and pyridyl; wherein R¹ is selected from 5- and 6-membered heterocyclo, lower cycloalkyl, lower cycloalkenyl and

aryl selected from phenyl, biphenyl and naphthyl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, cyano, carboxyl, lower alkoxy, lower alkoxyalkyl, lower hydroxyalkyl, lower haloalkoxy, amino, lower alkylamino, phenylamino, nitro, lower alkoxyalkyl, lower alkylsulfonate, halo, lower alkoxy and lower alkylthio; wherein R² is selected from lower alkyl and amino; and wherein R³ is a radical selected from halo, lower alkyl, oxo, cyano, carboxyl, lower cyanoalkyl, heteroaryloxy, lower alkyloxy, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower hydroxyalkyl, lower aralkyl, acyl, phenylcarbonyl, lower alkoxyalkyl, heteroaryloxy, alkoxy, alkoxyalkyl, aminocarbonyl, alkylaminocarbonyl, alkylamino, aminoalkyl, alkylaminoalkyl, aryloxy, and aralkoxy; or a pharmaceutically-acceptable salt thereof.

15. The combination of Claim 14 wherein A is selected from oxazolyl, isoxazolyl, dihydrofuryl, imidazolyl, and pyrazolyl; wherein R¹ is selected from 5- and 6-membered heterocyclo, and aryl selected from phenyl, biphenyl and naphthyl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, cyano, carboxyl, lower alkoxy carbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino, lower alkylamino, phenylamino, nitro, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio; wherein R² is amino; and wherein R³ is a radical selected from oxo, cyano, carboxyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, halo, lower alkyl, lower alkyloxy, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower hydroxylalkyl, lower aralkyl, acyl, phenylcarbonyl, lower alkoxyalkyl, 5- or 6-membered heteroaryloxy, aminocarbonyl, lower alkylaminocarbonyl,

lower alkylamino, lower aminoalkyl, lower alkylaminoalkyl, phenyloxy, and lower aralkoxy; or a pharmaceutically-acceptable salt thereof.

- 5 16. The combination of Claim 15 wherein A is
selected from oxazolyl, isoxazolyl, imidazolyl, and
pyrazolyl; wherein R¹ is phenyl optionally substituted
at a substitutable position with one or more radicals
selected from methyl, ethyl, isopropyl, butyl, tert-
10 butyl, isobutyl, pentyl, hexyl, trifluoromethyl, cyano,
carboxyl, methoxycarbonyl, hydroxyl, hydroxymethyl,
trifluoromethoxy, amino, N-methylamino, N,N-
dimethylamino, N-ethylamino, N,N-dipropylamino, N-
butylamino, N-methyl-N-ethylamino, nitro,
15 methoxymethyl, methylsulfinyl, fluoro, chloro, bromo,
methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and
methylthio; wherein R² is amino; and wherein R³ is a
radical selected from oxo, cyano, carboxyl,
methoxycarbonyl, ethoxycarbonyl, carboxypropyl,
20 carboxymethyl, carboxyethyl, cyanomethyl, fluoro,
chloro, bromo, methyl, ethyl, isopropyl, butyl, tert-
butyl, isobutyl, pentyl, hexyl, fluoromethyl,
difluoromethyl, trifluoromethyl, chloromethyl,
dichloromethyl, trichloromethyl, pentafluoroethyl,
25 heptafluoropropyl, fluoromethyl, difluoroethyl,
difluoropropyl, dichloroethyl, dichloropropyl, methoxy,
ethoxy, propoxy, n-butoxy, pentoxy, cyclohexyl, phenyl,
pyridyl, thienyl, thiazolyl, oxazolyl, furyl,
pyrazinyl, hydroxylmethyl, hydroxylpropyl, benzyl,
30 formyl, phenylcarbonyl, methoxymethyl, furylmethoxy,
aminocarbonyl, N-methylaminocarbonyl, N,N-
dimethylaminocarbonyl, N,N-dimethylamino, N-ethylamino,
N,N-dipropylamino, N-butylamino, N-methyl-N-ethylamino,
aminomethyl, N,N-dimethylaminomethyl, N-methyl-N-
35 ethylaminomethyl, benzyloxy, and phenyloxy; or a
pharmaceutically-acceptable salt thereof.

17. The combination of Claim 16 wherein the cyclooxygenase-2 inhibitor is selected from compounds, their prodrugs and their pharmaceutically-acceptable salts, of the group consisting of

- 5
- 3-(3,4-difluorophenyl)-4-(4-methylsulfonylphenyl)-2-(5H)-furanone;
- 3-phenyl-4-(4-methylsulfonylphenyl)-2-(5H)-furanone;
- 10 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 15 3-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;
- 2-methyl-5-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;
- 4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;
- 20 4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide;
- 4-[5-hydroxyethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;
- [2-trifluoromethyl-5-(3,4-difluorophenyl)-4-oxazolyl]benzenesulfonamide;
- 25 4-[2-methyl-4-phenyl-5-oxazolyl]benzenesulfonamide; and
- 4-[5-(3-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-oxazolyl]benzenesulfonamide.

- 30 18. The composition of Claim 10 wherein the leukocyte activation inhibitor is a cyclosporin.


19. The composition of Claim 18 wherein the cyclosporin is cyclosporin A.

35

20. A pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a therapeutically-effective amount of a leukotriene B₄

receptor antagonist, a cyclosporin and a cyclooxygenase-2 inhibitor selected from Dupont Dup 697, Taisho NS-398, meloxicam, flosulide and compounds of Formula I

5



wherein A is a 5- or 6-member ring substituent selected from partially unsaturated or unsaturated heterocyclo and carbocyclic rings;

wherein R¹ is at least one substituent selected from heterocyclo, cycloalkyl, cycloalkenyl and aryl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxy carbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

wherein R² is selected from alkyl, and amino; and

wherein R³ is a radical selected from halo, alkyl, alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl, heterocycloxy, alkyl, alkoxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, heterocyclo, cycloalkenyl, aralkyl, heterocycloalkyl, acyl, alkylthioalkyl, hydroxyalkyl, alkoxy carbonyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl, aryloxyalkyl, aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl, alkoxy carbonylalkyl, aminocarbonyl, aminocarbonylalkyl, alkylaminocarbonyl, N-arylaminocarbonyl, N-alkyl-N-arylaminocarbonyl, alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-aryl amino, N-aralkylamino, N-alkyl-N-aralkylamino, N-alkyl-N-aryl amino, aminoalkyl, alkylaminoalkyl, N-aryl aminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-aralkylaminoalkyl, N-alkyl-N-aryl aminoalkyl, aryloxy,

aralkoxy, arylthio, aralkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-arylamino sulfonyl, arylsulfonyl, N-alkyl-N-arylamino sulfonyl;

5 or a pharmaceutically-acceptable salt thereof.

21. The method of Claim 1 further characterized by suppressing immune response in a subject susceptible to or afflicted with rejection of an organ transplanted to said subject; an autoimmune disease, an inflammatory disease, or a condition with underlying autoimmune or inflammatory reactivities or responses; a graft versus host disease; an allergy; asthma; airway hypersensitivity; septic shock; myasthenia gravis; 15 autoimmune thyroiditis; Grave's disease; autoimmune hemolytic anemia; autoimmune thrombocytopenia purpura; mixed connective tissue disease; idiopathic Addison's disease; Sjogren's syndrome; urticaria; an acute hypersensitivity response or a delayed hypersensitivity response; Goodpasture's syndrome; hemolytic anemia; 20 contact dermatitis; granuloma; antibody-induced thrombocytopenia; hypersensitivity pneumonitis; glomerulonephritis; thyroiditis; encephalomyelitis; or meningitis.

25

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/01422

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K45/06 A61K31/00 A61K31/10 A61K31/18 A61K38/13

According to International Patent Classification (IPC) or in both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 96 41645 A (G.D.SEARLE & CO.) 27 December 1996 see the whole document ---	10-17,21
X	DE 42 28 201 A (SCHERING AG.) 3 March 1994 see the whole document see page 6, line 55 - line 59 ---	1-21
Y	WO 96 03385 A (SEARLE G.D. & CO.) 8 February 1996 cited in the application see the whole document ---	1-21
Y	WO 95 15316 A (SEARLE G.D. & CO.) 8 June 1995 cited in the application see the whole document ---	1-21
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

11 July 1997

Date of mailing of the international search report

21.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Economou, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/01422

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 03388 A (SEARLE G.D. & CO.) 8 February 1996 cited in the application see the whole document ---	1-21
Y	WO 96 03387 A (SEARLE G.D. & CO.) 8 February 1996 cited in the application see the whole document ---	1-21
Y	US 5 384 318 A (PFIZER INC.) 24 January 1995 cited in the application see the whole document ---	1-21
Y	US 5 246 965 A (CIBA-GEIGY) 21 September 1993 cited in the application see the whole document ---	1-21
A	EXPERT OPINION OF INVESTIGATIONAL DRUGS, vol. 4, no. 9, 16 - 19 June 1995, U.K., pages 885-888, XP000603401 PARNHAM M.: "PULMONARY -ALLERGY, DERMATOLOGICAL, GASTROINTESTINAL & ARTHRITIS" see page 888, right-hand column, last paragraph - page 887, right-hand column, line 5 ---	1-21
P,Y	DRUG NEWS AND PERSPECTIVES, vol. 9, no. 10, 27 - 31 October 1996, SPAIN, pages 631-639, XP000655227 PARNHAM M.: "INFLAMMATION:MECHANISMS AND THERAPEUTICS" see page 635, right-hand column, line 15 - page 638, middle column, line 24 ---	1-21
Y	EXPERT OPIN. INVEST. DRUGS, vol. 3, no. 2, 1994, U.K., pages 185-190, XP000610416 BROOKS D.W.: "PROGRESS WITH INVESTIGATIONAL DRUGS FOR THE TREATMENT OF PULMONARY AND INFLAMMATORY DISEASES" see the whole document ---	1-21
Y	EXPERT OPINION ON INVESTIGATIONAL DRUGS, vol. 4, no. 10, 1995, U.K., pages 1021-1025, XP000603399 CARRUTHERS N.I. ET AL.: "RECENT DEVELOPMENTS WITH INVESTIGATIONAL DRUGS POTENTIALLY USEFULL IN THE TREATMENT OF ALLERGIC AND INFLAMMATORY DISORDERS" see the whole document -----	1-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/01422

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9641645 A	27-12-96	AU 6269496 A	09-01-97
DE 4228201 A	03-03-94	AU 4950593 A	15-03-94
		CA 2142821 A	03-03-94
		WO 9404522 A	03-03-94
		EP 0656896 A	14-06-95
		HU 70511 A	30-10-95
		JP 8500585 T	23-01-96
		NO 950708 A	24-02-95
		US 5559134 A	24-09-96
WO 9603385 A	08-02-96	US 5486534 A	23-01-96
		AU 3126795 A	22-02-96
		CA 2195123 A	08-02-96
		EP 0772597 A	14-05-97
		US 5580985 A	03-12-96
WO 9515316 A	08-06-95	US 5466823 A	14-11-95
		US 5521207 A	28-05-96
		AU 1171495 A	19-06-95
		CA 2177576 A	08-06-95
		CN 1141630 A	29-01-97
		CZ 9601503 A	11-12-96
		EP 0731795 A	18-09-96
		FI 962249 A	29-05-96
		HU 74180 A	28-11-96
		NO 962184 A	29-05-96
		PL 314695 A	16-09-96
		US 5510496 A	23-04-96
		US 5563165 A	08-10-96
		US 5508426 A	16-04-96
		US 5516907 A	14-05-96
		US 5504215 A	02-04-96
		ZA 9409418 A	28-11-95
WO 9603388 A	08-02-96	US 5616601 A	01-04-97
		AU 3202595 A	22-02-96
		CA 2195845 A	08-02-96
		EP 0772600 A	14-05-97

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/01422

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9603387 A	08-02-96	US 5620999 A	15-04-97
		AU 3271695 A	22-02-96
		CA 2195846 A	08-02-96
		EP 0772601 A	14-05-97

US 5384318 A	24-01-95	WO 9015801 A	27-12-90
		CA 2019349 A	22-12-90
		EP 0404440 A	27-12-90
		FI 96951 B	14-06-96
		JP 3038569 A	19-02-91
		JP 7053722 B	07-06-95

US 5246965 A	21-09-93	AU 1807292 A	17-12-92
		CA 2070796 A	12-12-92
		CS 9201781 A	16-12-92
		EP 0518818 A	16-12-92
		HU 61977 A	29-03-93
		JP 5239008 A	17-09-93
